

1-11-2010

Candida species in cystic fibrosis: A road less travelled.

Sanjay H. Chotirmall

Royal College of Surgeons in Ireland

Catherine M. Greene

Royal College of Surgeons in Ireland, cmgreene@rcsi.ie

Noel G. McElvaney

Royal College of Surgeons in Ireland

Citation

Chotirmall SH, Greene CM, McElvaney NG. Candida species in cystic fibrosis: A road less travelled. *Medical Mycology*. 2010;48 Suppl 1:S114-24.

This Article is brought to you for free and open access by the Department of Medicine at e-publications@RCSI. It has been accepted for inclusion in Medicine Articles by an authorized administrator of e-publications@RCSI. For more information, please contact epubs@rcsi.ie.

Attribution-Non-Commercial-ShareAlike 1.0

You are free:

- to copy, distribute, display, and perform the work.
- to make derivative works.

Under the following conditions:

- Attribution — You must give the original author credit.
- Non-Commercial — You may not use this work for commercial purposes.
- Share Alike — If you alter, transform, or build upon this work, you may distribute the resulting work only under a licence identical to this one.

For any reuse or distribution, you must make clear to others the licence terms of this work. Any of these conditions can be waived if you get permission from the author.

Your fair use and other rights are in no way affected by the above.

This work is licenced under the Creative Commons Attribution-Non-Commercial-ShareAlike License. To view a copy of this licence, visit:

URL (human-readable summary):

- <http://creativecommons.org/licenses/by-nc-sa/1.0/>

URL (legal code):

- <http://creativecommons.org/worldwide/uk/translated-license>
-



CANDIDA SPECIES IN CYSTIC FIBROSIS: A ROAD LESS TRAVELLED

Journal:	<i>Medical Mycology</i>
Manuscript ID:	Draft
Manuscript Type:	Review Article
Date Submitted by the Author:	
Complete List of Authors:	Chotirmall, Sanjay; Royal College of Surgeons in Ireland, Respiratory Research Division, Department of Medicine Greene, Catherine; Royal College of Surgeons in Ireland, Respiratory Research Division, Department of Medicine McElvaney, Noel; Royal College of Surgeons in Ireland, Respiratory Research Division, Department of Medicine
Keyword:	Candida, Infection , Colonization , Immune response , Cystic Fibrosis



CANDIDA SPECIES IN CYSTIC FIBROSIS: A ROAD LESS TRAVELLED

**#Sanjay H. Chotirmall, *Catherine M. Greene & Noel G. McElvaney*

*These authors contributed equally to the work (Joint 1st authors)

#Member of the ISHAM working group for fungal infection in cystic fibrosis

Respiratory Research Division, Department of Medicine,
Royal College of Surgeons in Ireland

Corresponding Author:

Dr Sanjay H. Chotirmall

Respiratory Research Division, Education & Research Centre,

Beaumont Hospital, Beaumont Road, Dublin 9, Republic of Ireland

E-mail – schotirmall@rcsi.ie

Contact Number – 353-87-979-3833

Fax Number – 353-1-8093765

Running title: The role of *Candida spp* in CF

Keywords: *Candida*, Infection, Colonization, Immune response, Cystic Fibrosis

Word Count (Text): 5,925, Word Count (Abstract): 162

Disclosure: The authors report no conflicts of interest.

Abstract

Candida species are isolated with high frequency in cystic fibrosis yet their definitive role in disease remains unclear. Previously considered to have minimal inherent virulence owing to their commensal ability, the last decade has heralded an increasing recognition of *Candida* infection among patients with cystic fibrosis. What has been more recently hypothesized is that the organism possesses virulence factors that play diverse roles at different body sites during varied stages of an infection. Currently, limited data is accessible in the area of cystic fibrosis. This review aims to provide an overview of the role of *Candida* species in cystic fibrosis as is currently understood including the common local and systemic infections observed in clinical practice. The uncertain role of airway colonization and insight into emerging fields such as *Candida*-bacterial interactions are also addressed. Finally, we outline the current understanding of the innate, cellular and humoral immune responses associated with this genus which has been the major focus of work performed to date.

1. Introduction

Major advances in the care of cystic fibrosis (CF) patients have positively influenced prognosis over the last decade. Significant inroads into understanding the basic defect have accelerated the development of targeted therapies. The disease however continues to present new challenges to clinicians and researchers, for example fungal airway colonization. The consequences of bacterial infection, colonization and need for segregation in outpatient clinics have all been confronted and curbed to the point that fungal colonizers with an undetermined role on disease course and progression are becoming increasingly prevalent. Both yeasts and filamentous fungi have been identified as microbial pathogens in CF particularly in the context of invasive disease in the transplanted population and allergic responses, for instance allergic bronchopulmonary aspergillosis (ABPA). One particular fungal genus isolated at high frequencies from sputum culture is *Candida* and limited literature is available addressing the issues of *Candida* colonization and infection in CF. This is possibly because its manifestations are still considered relatively minor in comparison to other infectious agents. As a consequence it has received little attention in terms of clinical and scientific research. This review aims to provide an overview and our current understanding of the *Candida* species in CF, its associated local and systemic infections and an insight into emerging data on its role in airway colonization and associated immune response.

2. The *Candida* genus

Oral thrush was the first infection of the *Candida* species described in humans and following identification of its reproductive potential by budding, the fungus was originally named *Oidium albicans*. *Candida albicans* became the adopted name used and since then many other species have been identified to play a role in human infection. The most common of

1
2
3 these are *Candida albicans*, *Candida glabrata*, *Candida krusei*, *Candida parapsilosis* and
4
5
6 *Candida tropicalis* (1, 2).
7
8
9

10 The genus has exponentially grown with hundreds of newer member species identified and
11 unlike dimorphic fungi the morphology of a given *Candida* species remains fundamentally
12 comparable *in vitro* or *vivo*. *Candida* species are capable of causing chronic, localized or
13 systemic infection collectively termed 'candidiasis' although most commonly act as
14 commensals within the oropharynx, skin folds, gastrointestinal tract and vagina. On occasion
15 it can cause opportunistic infection (3). Once infection ensues, significant morbidity and
16 mortality results and consequently, systemic candidiasis has high death rates (>75%) (4).
17
18
19
20
21
22
23
24
25
26
27
28

29 Identification in the microbiology laboratory is achieved on Sabouraud dextrose media and
30
31 *Candida albicans* can be identified through germ tube testing or colorimetric detection of L-
32 proline aminopeptidase and beta-galactosaminidase. Since recent isolation of *Candida*
33 *dubliniensis* which produces false positive results in the above tests, a chromogenic agar
34 culture method that allows isolation and identification of *Candida albicans*, *Candida*
35 *tropicalis* and *Candida krusei* has been widely adopted. A modified version of this media is
36 available to more clearly distinguish *Candida dubliniensis* (5). Alternative older methods to
37 distinguish between the two organisms include an assessment of growth ability at higher
38 temperatures (45°C for *Candida albicans*) or specific DNA sequencing.
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 Patients with CF are at an increased risk of acquiring *Candida* due to use of inhaled steroids,
54 diabetes mellitus and lifelong antibiotic treatment however despite its frequent isolation from
55 sputum, oral and vaginal swabs, it remains unclear what such culture actually means in
56 practical terms for CF clinicians. We believe that a spectrum of "commensal-colonizer-
57
58
59
60

1
2
3 pathogen” most likely exists for the organism and where specifically the organism is on this
4
5 spectrum at a particular time point may be dictated by the clinical state of the CF patient and
6
7 whether bacterial co-colonizers are concurrently present in the airway.
8
9

10
11
12 Prior studies have addressed the notion that although frequently identified in CF, the clinical
13
14 role of *Candida* species has yet to be definitively determined (6-8). *Bakare* et al identified
15
16 *Candida* as the second most frequent fungal growth to *Aspergillus* in the CF airway and such
17
18 growth has been associated with more severe CF where patients receive prolonged treatment
19
20 with antibiotics, glucocorticoids and probiotics (9-12). In terms of infection, *Cimon* et al
21
22 performed a five-year epidemiological study assessing the frequency of bronchopulmonary
23
24 mycoses in a CF population and examined the aetiological role of individual fungal species in
25
26 disease. The filamentous fungi *Aspergillus* and *Scedosporium apiospermum* together with
27
28 *Candida* contributed the largest burden. Despite high isolation of *Candida* in CF, a single
29
30 case of candidiasis was observed and this low rate was attributed to the anti-fungal ability of
31
32 various bacterial colonizers in CF and whilst invasive airway infection is a rare event, extent
33
34 of airway damage from hypersensitivity phenomena remain unknown (13). We will now
35
36 address the localized and systemic infections associated with *Candida* species in CF and
37
38 subsequently tackle the complex issues of airway colonization, cross kingdom interaction and
39
40 the immune response.
41
42
43
44
45
46
47
48
49
50

51 **3. Localized *Candida* infection in CF**

52 **3.1 Oral candidiasis**

53
54 *Candida* species are isolated from the oral mucosa in up to 40% of healthy adults and
55
56 therefore considered commensal (14). A cut-off point to distinguish between commensalism
57
58 and colonization remains undetermined. Common risk factors associated with oral recovery
59
60

1
2
3 include poor dentition, older age, diabetes mellitus, use of inhaled or systemic steroids,
4
5 smoking, malignancy and frequent antibiotic use. Oral thrush usually presents as discomfort
6
7 associated with a dry mouth and associated dysphagia. In some cases, altered taste is
8
9 experienced. The diagnosis is usually straightforward and by direct observation of white
10
11 membranous plaques on the buccal mucosa or soft palate. This may be confirmed
12
13 microbiologically by staining a swab or culturing a rinse from the associated area. Atypically,
14
15 foci of oral erythematous inflammation or angular cheilitis may present. There are clearly
16
17 significant risk factors in the CF state that predispose to oral colonization and subsequent
18
19 infection including impaired salivary secretion, steroid use, CF-related diabetes and recurrent
20
21 courses of antibiotics for exacerbations. Antibiotics alter the homeostasis of oral flora and as
22
23 such have a permissive action on *Candida* growth. In a study from Manchester, on direct
24
25 questioning of major symptoms in the CF population, 40% (n=17) complained of a sore
26
27 mouth, 24% (n=10) of thrush five times annually and 38% (n=16) of a hoarse voice every
28
29 three months (15). In our own institution's experience, we encounter regular instances of oral
30
31 candidiasis annually following courses of antibiotics but which resolve after a short burst of
32
33 anti-fungal treatment (Fluconazole). We recommend microbiological confirmation by
34
35 scrapings in all cases unless white plaques are directly observed on oral examination. This is
36
37 because some of the symptoms described are not specific to oral thrush but can be found in
38
39 associated vitamin deficiencies (B₆, B₁₂) or by simple blistering. We recommend that CF
40
41 patients attending routine clinic be screened for risk factors and questioned at three-monthly
42
43 intervals with regard to the symptoms of oral thrush including frequency of sore or dry
44
45 mouth, crusting lips, dysphagia, dysphonia or hoarseness and difficulties with taste. Any
46
47 relationship to antibiotic treatment should be teased out. Clearly oral candidiasis is
48
49 recognized in CF but the dearth of available literature suggests it is probably misdiagnosed,
50
51 ignored or missed in several cases. A simple risk factor and symptom screen would see
52
53
54
55
56
57
58
59
60

1
2
3 improvements both in quality of life and probably compliance with other CF treatments. With
4 the advent of newer and increasingly earlier administration of anti-bacterial therapies, oral
5 candidiasis is likely to become a more significant issue in the future care of CF patients.
6
7
8
9

10 11 12 13 **3.2 Genital candidiasis**

14
15 Genital candidiasis is a common occurrence in the normal population with rates of up to 75%
16 having single and 50% recurrent episodes (2). It may be asymptomatic or present with
17 balanitis in males and pruritis with vaginal discharge in females. Most infections are caused
18 by *Candida albicans* (95%) however *Candida glabrata* (5%) infection is described (2). There
19 is limited but important literature available addressing these infections in CF with the
20 majority focused on female manifestations. It has been more than a decade since *Sawyer et al*
21 first reviewed the subject with a self-administered questionnaire in young women with CF
22 (n=55) (16). Vulvovaginal candidiasis was more common in CF (35%) versus controls (13%)
23 and additionally more persistent and difficult to treat. Antibiotic use was a significant
24 association and the work concluded that “health professionals generally trivialize illnesses
25 and diseases that are common, easily treated and not life-threatening”. More recent work has
26 included male patients and addressed symptomatic partners. *Lyon et al* evaluated 40 adults
27 with CF (19 male, 21 female) and similarly found large proportions (62.5%, n=25)
28 experiencing symptoms of infection however few (15%, n=6) had been directly questioned
29 about it at CF clinic (17). Patients refused to discuss if their partners were symptomatic. This
30 highlighted to CF clinicians a major deficiency of clinical practice and questions about
31 candidiasis should feature during annual review clinic consultations. It is also important to
32 consider that vaginal discharge in young CF women can be caused by other pathogens such
33 as *Chylamidia*, *Gonococcus* or *Trichomonas* species and that there is an observed
34 unexplained high incidence of genital *Chylamidia* in CF.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A third study addressing the same subject was performed by interview in 101 CF patients and
4 addressed symptom frequency and medical risks associated with *Candida* (18). Patients were
5 asked to report on personal risk factors for *Candida* infection and their desire to be
6 questioned about 'thrush' in CF clinic. Many had two or more risk factors (92.1%, n=93)
7 however the only significant factor associated with genital *Candida* was long-term antibiotics
8 (87.1%, n=88, p=0.001). Over seventy percent of patients, both males and females had
9 symptoms of either oral or genital *Candida* or both simultaneously. Forty percent (18/45) of
10 those with oral and two-thirds (33/50) of those reporting genital candidiasis described
11 'distress' however it did not affect desire for treatment. Most cases of oral infection were
12 diagnosed by a CF physician whilst genital infection was mainly self-diagnosed. It is
13 noteworthy that general practitioners diagnosed more cases of genital infection when
14 compared to CF physicians. This is potentially explained by differing doctor-patient
15 relationships in different settings or alternatively because the focus of the CF unit remains on
16 respiratory or gastrointestinal symptoms leading patients to believe that this forum is
17 inappropriate for discussing other complaints. Most patients in the study did however want to
18 discuss such issues and were unconcerned who their discussant was although some females
19 predictably preferred discussion with female staff. This Manchester based study is the largest
20 to date and detected similar patterns to previous work. The most concerning new discovery
21 was the high incidence of symptoms among CF patients but only on direct questioning
22 placing a future onus on CF clinics. A major criticism of all these studies was that symptom
23 recording was not supported by microbiological confirmation of infection, an important point
24 for future work. Additionally, although several publications have assessed the benefits and
25 efficacy of anti-fungal treatment in vulvovaginal candidiasis in the 'normal' population,
26 importantly none have been performed in CF (19). Despite this clear lack of available
27 literature, we strongly recommend screening questions for infection at all CF clinic visits and
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 depending on clinical findings, anti-fungal treatment prescribed either empirically or
4 following microbiological confirmation.
5
6
7
8
9

10 **4. Systemic *Candida* infection in CF**

11 **4.1 Post-transplant Candidiasis**

12
13 CF is the 3rd most common indication for lung transplantation and the opportunistic nature of
14
15 *Candida* species suggest that the post-transplant period is ripe for such infection (20). Despite
16
17 this, candidiasis post-transplant remains rare and *Aspergillus* species are in fact more
18
19 commonly encountered in this setting (21). The main *Candida* infection following
20
21 transplantation is surprisingly tracheobronchitis which includes anastomotic site infections.
22
23 Bloodstream and other invasive infections secondary to *Candida* are rare and will not be
24
25 addressed in any detail within this review except to state that when present occur within the
26
27 first month following transplantation (22, 23). This is primarily a consequence of the major
28
29 surgical intervention and intensive care unit stay experienced by patients.
30
31
32
33
34
35
36
37
38

39 **4.2 Totally implantable venous access device (TIVAD) infection**

40
41 A more commonly encountered systemic infection associated with *Candida* involves the
42
43 presence of a TIVAD commonly referred to as a “port”. *Candida* species in this setting are
44
45 recognized as the most common infecting organism associated with a TIVAD resulting in
46
47 septicaemia (24-26). Important risk factors for infection remain the same as that for other
48
49 *Candida* infections. Diagnosed by the presence of swinging pyrexia, systemic septicaemia
50
51 and positive blood cultures for *Candida* species taken from both the port site and
52
53 peripherally, the first intervention remains to remove the offending device whose tip should
54
55 also be sent for microbiological evaluation. Device removal results in significant clinical
56
57 improvement however aggressive anti-fungal therapy is concurrently administered during
58
59
60

1
2
3 which time patients on the active transplant list have to be removed temporarily. Recently, we
4
5 encountered a case series at our centre which presented a different setting to the traditional
6
7 *Candida* port infection. We experienced three cases of TIVAD thrombosis and superior vena
8
9 cava obstruction that required use of thrombolytic therapy. In two of the three patients, their
10
11 post-thrombolysis course was complicated by systemic candidiasis secondary to TIVAD
12
13 infection. In these cases, we achieved a successful outcome following removal of the device
14
15 coupled with aggressive anti-fungal treatment. Another more traditional case series described
16
17 earlier this decade over a six year period was that from a CF centre in Manchester where
18
19 fifteen adults with CF were diagnosed via positive blood cultures with a *Candida* port
20
21 infection (15). Here, a variety of *Candida* isolates were identified including *albicans*,
22
23 *parapsilosis* and *glabrata* and excellent clinical outcomes again achieved via device removal
24
25 and systemic anti-fungal treatment dictated through sensitivity testing. Our own practice
26
27 continues to evolve with regard to optimal treatment and we routinely look for at least two
28
29 negative blood cultures following completion of the prescribed treatment course. Replacing
30
31 ports depends on need but we try not to replace before 8 weeks following the last negative
32
33 blood culture. To date, there remain no clinical trials or evidence based guidelines to support
34
35 these treatment practices. Another important point is that many *Candida* infections involve
36
37 biofilm formation particularly with indwelling vascular catheters in the context of CF. These
38
39 biofilms are microbiologically complex containing matrix enclosed microcolonies containing
40
41 yeasts and hyphae in a bilayer structure (27). Such *Candida* biofilms can be resistant to
42
43 conventional anti-fungals through a multitude of mechanisms and as such future research
44
45 needs to be conducted to determine the best and optimally standardized treatment of *Candida*
46
47 port infections.
48
49
50
51
52
53
54
55
56
57
58
59
60

5. Airway *Candida* colonization in CF

It remains controversial as to whether *Candida* species are transient or persistent colonizers of the respiratory tract in CF. A study by Muthig et al showed that the mean persistence of *Candida* species was at least nine months and that the species identified were genetically related and transmissible but susceptible to all anti-fungals tested. Although concerns of transmissibility persist, it is unsure whether this species conclusively contributes to chronic infection and the inflammatory milieu in CF (28). We have assessed colonization rates at our own centre over prolonged time periods and found persistence rates in excess to that previously described. We have established that the main factors predicting colonization by *Candida albicans* in CF are pancreatic insufficiency, osteopenia and co-colonization with *Pseudomonas*. At first glance, this suggests that the more advanced a patient's disease, the likelier their sputum contained *Candida albicans*, a view of many clinicians and it may be that the organism acts as nothing more than a microbiological marker of disease severity in CF. To challenge this paradigm, we are currently prospectively evaluating whether airway colonization by *Candida albicans* may act pathogenically by affecting clinical outcomes in CF including FEV1, BMI, hospitalizations for infective exacerbations and sputum colonization with *Pseudomonas* or *Aspergillus* species. Notably, a previous cross sectional analysis of a European CF registry did show that *Candida albicans* colonization was associated with 5-10% predicted decrease in pulmonary function (29).

Newer airway *Candida* species have also emerged over the last decade and one pertinent example is the high recovery rates (10-25%) of *C. dubliniensis* from the oral cavity of HIV patients (30). This new organism was subsequently described in the non-HIV population particularly in individuals receiving high antibiotic burdens (31). Therefore, its detection in CF came as no surprise but did involve a complex isolation procedure involving Staib agar

1
2
3 (32). What was surprising was that its prevalence rate in CF was higher than that found in
4
5 HIV however virtually nothing about its potential for virulence is known. There is no clear
6
7 clinical or experimental evidence of differences in terms of pathogenic potential when
8
9 compared to *Candida albicans* (33) however *Candida dubliniensis* is reported to exhibit cell
10
11 surface hydrophobicity not observed in the *albicans* species (34). Cell surface hydrophobicity
12
13 is known to play a role in the adhesion of microorganisms and by displaying this feature,
14
15 *Candida dubliniensis* takes advantage of the dehydrated respiratory secretions in CF and
16
17 consequently proliferates. These observations may also explain why patients who are older
18
19 (>30 years) and have more advanced disease are colonized by this yeast. *Peltroch et al* (32)
20
21 followed six CF patients with *Candida dubliniensis* and whilst all patients remained stable
22
23 with no invasive infection detected its effect on lung function could not be conclusively
24
25 established because of the small numbers. A larger epidemiological study of this *Candida*
26
27 species in CF is warranted.
28
29
30
31
32
33

34 35 36 **6. *Candida*-bacterial interactions** 37

38
39 According to Costerton (35), biofilms are 'a structured community of bacterial cells enclosed
40
41 in a self-produced polymeric matrix and adherent to an inert or living surface'. Traditionally
42
43 biofilms have been thought to comprise a single bacterial species however it is now
44
45 increasingly recognised that mixed biofilms exist involving interactions between both
46
47 prokaryotes and eukaryotes. Bacteria and fungi are found together in a variety of
48
49 environments but particularly in biofilms, where adherent species interact through diverse
50
51 signaling mechanisms. In the host *C. albicans* can often be found growing with bacteria in
52
53 polymicrobial biofilms and interspecies interactions occur that can impact on the transition of
54
55 *C. albicans* between virulent and nonvirulent states (27). Under conditions of immune
56
57 dysfunction, such as in the CF lung, colonising *C. albicans* can become an opportunistic
58
59
60

1
2
3 pathogen causing mucosal and disseminated infections potentially impacting on mortality. In
4
5 the biofilm environment, microbial species use 'quorum-sensing' (QS) molecules for cell-to-
6
7 cell communication to promote collective behaviour within the population, enhance access to
8
9 nutrients and niches, and provide a combined defense against competitor organisms (36, 37).
10
11 The process of QS can cross the prokaryote–eukaryote boundary (36-39).
12
13
14
15
16
17

18 **6.1 Interactions with *Pseudomonas aeruginosa***

19
20 *Pseudomonas aeruginosa* is the most prevalent opportunistic pathogen in individuals with CF
21
22 and is the principal organism associated with biofilm formation in the CF lung; *S. aureus* and
23
24 *Burkholderia* spp. are also considered important pulmonary pathogens in CF. The dimorphic
25
26 yeast *C. albicans* is the most common eukaryotic microbe isolated from CF patient sputum (9,
27
28 10, 40). *C. albicans* can exist in a mixed biofilm where the prokaryotic and eukaryotic
29
30 communities exhibit either synergistic or antagonistic interactions. Several studies suggest
31
32 that *P. aeruginosa* and *C. albicans* interact with each other *in vivo*, and they are commonly
33
34 found together in mixed infections (41). In their seminal paper Hogan and Kolter (38) first
35
36 reported a pathogenic relationship between *P. aeruginosa* and *C. albicans*. They
37
38 demonstrated how *P. aeruginosa* can form a dense biofilm on *C. albicans* filaments and kill
39
40 the fungus. Interestingly this only occurred when *C. albicans* was growing in its filamentous
41
42 (or hyphal) form – an essential feature associated with its virulence (42). *P. aeruginosa*
43
44 neither bound to nor killed the yeast form of *C. albicans* and the ability of *P. aeruginosa* to
45
46 kill filamentous *C. albicans* was dependent on a number of physiological factors including
47
48 growth phase, nutrient availability, surface structures including flagellae and type IV pili,
49
50 secreted QS factors and regulatory molecules such as *rpoN* (43). By forming a biofilm on
51
52 fungal filaments *P. aeruginosa* may be able to obtain nutrients from *C. albicans* in a
53
54 nutritionally scarce environment.
55
56
57
58
59
60

1
2
3 Prior to killing of *C. albicans* by *P. aeruginosa*, signalling can occur between both
4
5 organisms. The QS molecules of both species are responsible for this communication. For
6
7 example the bacterial molecule 3-oxo-C12 homoserine lactone can affect *Candida*
8
9 morphology, whilst the fungal 12-carbon sesquiterpene metabolite, farnesol can interfere with
10
11 *Pseudomonas* quinolone and pyocyanin production and swarming motility (37, 39, 41, 44-
12
13 49). Thus eukaryotes and prokaryotes possess diverse signaling mechanisms to detect and
14
15 respond to each other through QS signal molecules.
16
17
18
19
20

21 **6.2 Interactions with *Staphylococcus aureus***

22
23 In oral biofilms a mutually beneficial interaction called coaggregation can occur where the
24
25 adhesion of *C. albicans* to oral bacteria facilitates its colonization of the oral cavity (50-52).
26
27 In contrast, the interaction between *C. albicans* and *Pseudomonas aeruginosa* as described
28
29 above is competitive and antagonistic in nature. A third mechanism of interaction that can
30
31 occur is that evident between staphylococci and *C. albicans*, which appears to be initially
32
33 synergistic (53-55). Carlson *et al.* (56, 57) described a synergistic effect between *C. albicans*
34
35 and *S. aureus* in a mouse infection model leading to enhanced mortality following dual
36
37 infection suggesting that *C. albicans* can either enhance the virulence of *S. aureus* or impair
38
39 the host's immune defences. Extensive physical interactions are known to occur between *S.*
40
41 *aureus* and both the yeast and hyphal forms of *C. albicans* in a mixed biofilm (58) and it has
42
43 been suggested that farnesol has a role in orchestrating these interactions. After the initial
44
45 synergy during *C. albicans-S. aureus* biofilm formation farnesol then negatively affects
46
47 staphylococcal biofilm formation, compromises cell membrane integrity, viability and
48
49 susceptibility of *S. aureus* to a variety of clinically important antibiotics (58). Thus farnesol
50
51 may represent a therapeutic target for inhibiting the development of a mixed biofilm in the
52
53 CF lung however once the biofilm has been established farnesol may actually behave as an
54
55
56
57
58
59
60

1
2
3 anti-bacterial factor. It remains to be seen which has the more detrimental effect in the CF
4 lung, *C. albicans* growing alone or in combination with *S. aureus* in a mixed biofilm.
5
6
7
8

9 **6.3 Interactions with other microbes**

10
11 Notwithstanding the ability of *C. albicans* to modulate bacterial growth, reciprocal evidence
12 indicates that other bacteria may also play an important role in the pathogenesis of *C.*
13 *albicans* infections. For example in the urinary tract *Escherichia coli* can enhance adhesion of
14 *C. albicans* to bladder mucosa (59) whereas in the gut indigenous microbes can inhibit
15 mucosal adhesion of *C. albicans* (60). Consequently alterations in the normal bacterial flora
16 following treatment with broad-spectrum antibiotics may allow *C. albicans* to proliferate and
17 invade tissues, greatly affecting its pathogenicity (60). This is an important consideration for
18 individuals with CF who are frequently prescribed antibiotics.
19
20
21
22
23
24
25
26
27
28
29
30

31
32 This may be most clearly studied in the oral cavity where adhesion of *C. albicans* to saliva-
33 coated surfaces and proline-rich proteins is an important early step in colonization (50, 61-
34 63). Many species of oral bacteria may compete with *C. albicans* for primary adhesion
35 receptor sites (50, 61, 64, 65) however once resident in the mouth *C. albicans* can adhere to
36 the major microbial constituents of early dental plaque.
37
38
39
40
41
42
43

44
45 Similar to *P. aeruginosa*, another opportunistic pathogen *Acinetobacter baumannii* exhibits a
46 predilection for *C. albicans* filaments and can inhibit *C. albicans* filamentation, resulting in
47 attenuated virulence of *C. albicans* in the nematode. Interestingly, similar to its effect on *S.*
48 *aureus*, *C. albicans* can also inhibit *A. baumannii* growth via farnesol production (66). As
49 mixed bacterial–fungal biofilms have been shown to be associated with a multitude of
50 infections including those affecting endotracheal tubes, biliary stents, silicone voice and
51 orthopedic prostheses and acrylic dentures (35, 67) determining the exact sequence of events
52
53
54
55
56
57
58
59
60

1
2
3 involved in the development of these mixed biofilms may determine how and when to target
4
5 the individual microbial or fungal constituents.
6
7
8
9

10 11 **7. *Candida* and the immune response**

12
13 The outer strata of *Candida* species contain elements with antigenic potential. These include
14
15 mannans and mannoproteins which upon human exposure induce an immunogenic response
16
17 (68-71). Where mannan-deficient, *Candida* strains are clinically less virulent and during the
18
19 course of a *Candida* infection, cellular, humoral and innate immune responses all play a role
20
21 (69, 70, 72-74).
22
23
24
25
26
27

28 **7.1 Innate immunity**

29
30
31 Recognition of microbes by the innate immune system depends on activation of specific
32
33 pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRRs).
34
35 For fungi the first PAMPs encountered by the immune system are those present in the fungal
36
37 cell wall. The cell wall of *Candida albicans* is composed of a core structure of β -(1,3)-glucan
38
39 polysaccharide fibrils covalently linked to chitin (a β -(1,4)-linked polymer of *N*-acetyl
40
41 glucosamine) and β -(1,6)-glucan. The outer layer consists of *N*-linked (75) or *O*-linked
42
43 mannosylated proteins called mannans (76). Two classes of PRRs in particular play an
44
45 important role in antifungal immunity - the C-type lectin receptors (CLRs) and Toll-like
46
47 receptors (TLRs). Neutrophils, monocytes, macrophages and airway epithelial cells are all
48
49 involved in defense against fungal pathogens. Dendritic cells (DCs) also respond to fungal
50
51 PAMPs leading to activation of T-cell-mediated specific immunity. These various cell
52
53 populations differ in their expression of CLRs and TLRs on the cell membrane, and are
54
55 therefore capable of initiating different responses. CLRs and TLRs recognize the major
56
57
58
59
60

1
2
3 polysaccharide cell wall components, *N*- and *O*-linked mannans, β -mannosides, β -(1,6)-
4 glucan and phospholipomannan. The mannan structures are detected by the mannose
5 receptor (MR), the dendritic cell-specific ICAM3-grabbing nonintegrin (DC-SIGN), dectin-2,
6 galectin-3 and TLR4 whereas complement receptor 3 (CR3), dectin-1 and TLR2 detect the β -
7 glucans.
8
9
10
11
12
13
14

15 16 **7.1.1 CLRs** 17

18
19
20 CLRs comprise a large family of receptors that share at least one carbohydrate
21 recognition domain originally identified in mannose binding lectin (MBL). CLRs are
22 evolutionary conserved and have been shown to be involved in the modulation of the innate
23 immune response and fungal recognition. Although MBL can bind to *C. albicans* (77) and
24 has the ability to opsonize fungal yeasts by activating the complement system (78), MBL-
25 deficient mice do not show decreased survival to infection with *C. albicans* (79). However
26 dectins-1 and -2, galectin-3, DC-SIGN and MR do play important roles in the innate immune
27 response to *C. albicans*.
28
29
30
31
32
33
34
35
36
37
38
39

40 Dectin-1 recognizes β -(1,3)-glucans, mediates ligand uptake and phagocytosis, and
41 triggers cytokine production (80). Alone it is sufficient in inducing responses to fungi
42 however synergistic proinflammatory responses occur in cooperation with TLRs. For
43 example in collaboration with TLR2, dectin-1 triggers proinflammatory responses by *C.*
44 *albicans* or zymosan (81, 82). Dectin-2 is mainly expressed on myeloid cells and maturing
45 monocytes. It recognizes high-mannose structures (83) and interacts with the Fc γ R to induce
46 TNF in response to filamentous *C. albicans* (84). On macrophages the galectin-3 receptor
47 mediates the recognition β -mannosides expressed on *C. albicans*. This PAMP-PRR
48 interaction is also enhanced by TLR2 (85). DC-SIGN, expressed on mature DCs, recognizes
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 high-mannose structures in *C. albicans* and mediates phagocytosis of fungal particles (86).
4
5 Finally, MR recognizes chitin, fucose, and mannose and has been implicated in the
6
7 recognition of several fungi, including *C. neoformans*, *C. albicans*, and *Pneumocystis*.
8
9
10 Branched *N*-bound mannans in *C. albicans* are recognised by MR (87) and mice defective in
11
12 MR display partial impairment in their host defense against *Candida* infections (88).
13
14

15
16 Modulation of CLR expression and activity thus represent important therapeutic
17
18 targets in CF that remain, as yet, underexplored. A newly identified CLR, Mincle, has been
19
20 shown to participate in macrophage recognition of *C. albicans*. Although it remains to be
21
22 shown which PAMP expressed by *C. albicans* directly activates Mincle, the role of this
23
24 receptor in fungal innate immunity has been clearly demonstrated. Inhibition studies have
25
26 shown decreased TNF production by macrophages following stimulation by *Candida* yeast
27
28 cells and Mincle knockout mice display hypersusceptibility to *Candida* infection (89). It will
29
30 be interesting to determine the expression and function of Mincle by immune cells and
31
32 airway epithelium in individuals with CF.
33
34
35
36
37

38 39 **7.1.2 TLRs**

40
41
42 Following the initial observation by Lemaitre *et al.* (90) that *Drosophila melanogaster*
43
44 flies deficient in the Toll receptor succumb readily to infection with *Aspergillus fumigatus*
45
46 due to defective synthesis of the drosomycin defensin, the role for TLRs in antifungal defense
47
48 has been extensively studied and recently a human homolog of drosomycin called
49
50 drosomycin-like defensin has been described. This peptide is expressed mainly in the skin
51
52 and has activity against a variety of filamentous fungi (91), however its potential as an
53
54 antifungal or anti-*C. albicans* therapeutic has not yet been exploited. Early studies on TLRs
55
56 revealed that TLR2 and TLR6 co-operate in recognition of the fungal structure zymosan
57
58
59
60

1
2
3 derived from *Saccharomyces cerevisiae* (92). With respect to *C. albicans*, blocking of TLR2
4
5 has been shown to lead to decreased monocyte production of TNF and IL-1 β after stimulation
6
7
8 with *C. albicans* (93). Furthermore TLR2^{-/-} mice have decreased TNF and MIP-2 production
9
10 and reduced neutrophil recruitment after challenge with *Candida* (94). TLR1 and TLR6, two
11
12 receptors capable of forming heterodimers with TLR2, may also have a minor role in *C.*
13
14 *albicans* recognition (95). Further evidence for an anti-fungal role for TLR2 comes from
15
16 studies showing that TLR2-deficient macrophages have an increased ability to contain *C.*
17
18 *albicans* (96), and that TLR2 signalling can promote Th2-type or T-reg-type responses in
19
20 response to *C. albicans* (97, 98).
21
22
23
24
25

26 TLR4 participates in antifungal host defense by recognizing *O*-linked mannan
27
28 structures and mediating proinflammatory responses. TLR9 has the potential to recognize
29
30 fungal DNA and blocking TLR9 either pharmacologically in human monocytes or genetically
31
32 in TLR9-deficient mouse macrophages leads to a reduced production of cytokines, mainly
33
34 IL-10, in response to stimulation with *C. albicans* (99). However the contribution of TLR9 to
35
36 fungal recognition is not believed to be significant. Much is known regarding the expression
37
38 and function of TLRs in the CF lung and is beyond the scope of this article. Readers are
39
40 directly elsewhere for comprehensive reviews of the role of TLRs in CF (100, 101).
41
42
43
44
45

46 Recognition of *C. albicans* by the innate immune system therefore occurs through
47
48 MR and DC-SIGN recognizing branched *N*-linked mannans, and TLR4 recognizing linear *O*-
49
50 linked mannans. CR3 responds to β -(1,6)-glucan and dectin-1 and galectin-3 in combination
51
52 with TLR2 each recognize β -glucan/phospholipomannan and β -mannosides, respectively. It
53
54 is likely that these recognition receptors can operate in combination and that stimulation via
55
56 multiple PAMP-PRR combinations might increase both the sensitivity and the specificity of
57
58 the immune recognition process. Notwithstanding these elegant recognition systems, *C.*
59
60

1
2
3 *albicans* frequently colonizes individuals with CF. Exactly why *C. albicans* is so commonly
4
5 found in CF individuals remains to be determined but may be associated with impairment in a
6
7 particular component of the innate immune system.
8
9

10 11 **7.2 Cell mediated immunity**

12
13 Individuals whose cell mediated immunity is compromised are at an increased susceptibility
14
15 of infection with *Candida* species illustrating the important role that this arm of the immune
16
17 system plays in the host defence against the organism. The reasons underlying this
18
19 observation remain ambiguous and little work has been performed in CF to examine this
20
21 potential link. In non-CF animal models, interleukin (IL)-12 promotes a Th1 response to
22
23 *Candida* exposure and IL-10 systemic infection (102, 103). In contrast interferon (IFN)-
24
25 gamma appears to be protective (104). Allard et al have shown that CF oropharyngeal
26
27 exposure to *Candida* lysates evoked a Th2 type immune response similar to that observed in
28
29 non-CF models. However when exposed to both *Pseudomonas* and *Candida* lysates together
30
31 a deviation in the adaptive immune response from a Th2 to Th1 type associated with
32
33 neutrophilia is noted (105). Neutrophils also damage pseudohyphae in association with IFN-
34
35 gamma to provide another host resistance strategy against *Candida* species. This is mediated
36
37 through granulocyte colony stimulating factor (G-CSF) (106, 107). Additionally, Dectin-1 by
38
39 distinguishing cell wall β -glucan can trigger cell mediated defences (81). While the cytokine
40
41 milieu and neutrophil response in CF continue to be an area of intense investigation among
42
43 researchers, minimal data to date remains accessible with regard solely to *Candida* species.
44
45
46
47
48
49
50
51
52
53

54 **7.3 Humoral immunity**

55
56 The role of the humoral immune response during *Candida* exposure and infection remain
57
58 controversial. Despite this, the majority of literature with regards to *Candida* in CF exists in
59
60

1
2
3 this area. In the late 1980s, *Przyklenk et al* first assessed serum IgG antibodies to both
4
5 *Aspergillus fumigatus* and *Candida albicans* in CF versus control patients and found that
6
7 antibody levels were higher in CF irrespective of sputum isolation (108). In contrast to
8
9 *Aspergillus fumigatus*, antibodies to *Candida albicans* were observed to increase significantly
10
11 with increased isolation from sputum culture. Minimal work followed until *Maiz et al*
12
13 assessed the prevalence of *Aspergillus* and *Candida* species in CF sputa and the serologic IgE
14
15 responses to these fungi. For the first time, they additionally investigated whether the
16
17 immune response had direct effects on clinical status in CF (109). *Candida* species were
18
19 isolated in nearly half of all sputum samples analyzed (47.5%) however 87.9% of patients
20
21 had at least one growth of *Candida albicans* during the study course. One-quarter (26.7%)
22
23 were sensitized to *Candida albicans* and only patients who grew *Candida albicans* at least
24
25 once during the study developed an IgE response to the fungi. The clinical parameters
26
27 assessed (FEV1 and CT scores) were not worse in those sensitized versus the non-sensitized.
28
29 Interestingly, half of the sensitized group had confirmed ABPA whilst the remaining patients
30
31 some immunologic characteristics of ABPA. In conclusion, the group found a high
32
33 prevalence of both colonization and sensitization to *Candida albicans* in CF but could not
34
35 relate this to disease severity or clinical status. Although serum IgE to *Candida albicans*
36
37 appeared to represent an immunological marker of ABPA in CF, it is important to note that
38
39 the studied group was small (n=20) and only FEV1 and CT scores assessed as clinical
40
41 measures. The same group extended this work recently to assess serum IgG, IgA and IgM
42
43 against *Aspergillus fumigatus* and *Candida albicans* and found that although no correlation
44
45 was detected between the presence of *Aspergillus fumigatus* in sputum and an immune
46
47 response, the converse was true of *Candida albicans*. Increasing sputum isolation heralded an
48
49 elevated serum response however again this could not be related to respiratory impairment
50
51 (110).
52
53
54
55
56
57
58
59
60

8. Conclusion and future directions

In this review, we have highlighted the current knowledge base and infections caused by the *Candida* species in CF. The dearth of CF-specific literature available illustrates that it evidently is a 'road less travelled'. Despite this lack of literature and audit, what remains undoubted is that the species is isolated frequently and has importance in contributing to morbidity and in some cases mortality in CF. There is a high rate of undetected symptomatic oral and genital infection in the adult CF population and the problem should not be ignored with newer anti-bacterial agents on the horizon that will likely select out these fungal pathogens. Long-term use of antibiotics has recurrently emerged as a contributing factor to *Candida* infection and an alteration of flora post therapy lends survival advantages to the pathogenicity of this species. In doing so, *Candida* probably contributes to the inflammatory milieu observed in CF. Although post-transplant candidiasis is a rare occurrence, port infections do occur frequently. When a port is infected, it should be removed promptly and combined with anti-fungal therapy results in excellent clinical outcomes. The *Candida* species interestingly elicits innate, cellular and humoral immune responses that we have yet to fully understand in the context of CF. Clearly an increasing amount of work remains left to be done to address the many unanswered questions. Future avenues for focus in this field lie within clinical care, isolation techniques and biomedical research. Healthcare professionals should maintain a positive approach in looking for manifestations of *Candida* infection during annual review at CF clinics and subsequently pursue microbiology in symptomatic cases. In terms of isolation techniques, selective media needs to be developed to suppress the growth of gram negative pathogens such as *Pseudomonas* and *Burkholderia* species and enhance fungal identification and isolation. Standardization of detection protocols needs to be pursued for fungi in CF as currently lab and international variation persists. Finally, basic science and clinical research avenues with regard to *Candida* species in CF need to be

1
2
3 actively pursued so as to enable an improved understanding of its role in the CF airway,
4
5
6 *Candida*-bacterial interaction and its potential use as a microbiological marker of CF disease
7
8 severity and progression.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

References

1. Tavanti A, Davidson AD, Gow NA, Maiden MC, Odds FC. *Candida orthopsilosis* and *Candida metapsilosis* spp. nov. to replace *Candida parapsilosis* groups II and III. *J Clin Microbiol.* 2005 Jan;43(1):284-92.
2. D D. Fortnightly review: management of genital candidiasis. *BMJ.* 1995;10:1241-4.
3. Naglik JR, Challacombe SJ, Hube B. *Candida albicans* secreted aspartyl proteinases in virulence and pathogenesis. *Microbiol Mol Biol Rev.* 2003 Sep;67(3):400-28, table of contents.
4. Fraser VJ, Jones M, Dunkel J, Storfer S, Medoff G, Dunagan WC. Candidemia in a tertiary care hospital: epidemiology, risk factors, and predictors of mortality. *Clin Infect Dis.* 1992 Sep;15(3):414-21.
5. Sahand IH, Moragues MD, Eraso E, Villar-Vidal M, Quindos G, Ponton J. Supplementation of CHROMagar *Candida* medium with Pal's medium for rapid identification of *Candida dubliniensis*. *J Clin Microbiol.* 2005 Nov;43(11):5768-70.
6. Valenza G, Tappe D, Turnwald D, Frosch M, Konig C, Hebestreit H, et al. Prevalence and antimicrobial susceptibility of microorganisms isolated from sputa of patients with cystic fibrosis. *J Cyst Fibros.* 2008 Mar;7(2):123-7.
7. Kerem E, Conway S, Elborn S, Heijerman H. Standards of care for patients with cystic fibrosis: a European consensus. *J Cyst Fibros.* 2005 Mar;4(1):7-26.
8. Pihet M, Carrere J, Cimon B, Chabasse D, Delhaes L, Symoens F, et al. Occurrence and relevance of filamentous fungi in respiratory secretions of patients with cystic fibrosis--a review. *Med Mycol.* 2009 Jun;47(4):387-97.
9. Bakare N, Rickerts V, Bargon J, Just-Nubling G. Prevalence of *Aspergillus fumigatus* and other fungal species in the sputum of adult patients with cystic fibrosis. *Mycoses.* 2003 Feb;46(1-2):19-23.

10. Bauernfeind A, Bertele RM, Harms K, Horl G, Jungwirth R, Petermuller C, et al. Qualitative and quantitative microbiological analysis of sputa of 102 patients with cystic fibrosis. *Infection*. 1987 Jul-Aug;15(4):270-7.
11. Blaschke-Hellmessen R, Spitzer H, Paul KD, Hoffmann C. [Mycological surveillance of children with cystic fibrosis]. *Mycoses*. 1991;34 Suppl 1:43-7.
12. Muller J, Remus N, Harms KH. Mycoserological study of the treatment of paediatric cystic fibrosis patients with *Saccharomyces boulardii* (*Saccharomyces cerevisiae* Hansen CBS 5926). *Mycoses*. 1995 Mar-Apr;38(3-4):119-23.
13. Cimon B CJ, Chazalotte JP, Ginies JL, Chabasse D, Bouchara JP. Bronchopulmonary mycoses in cystic fibrosis. Results of a five-year longitudinal study. *Journal de Mycologie Medicale*. 2000;10(3):128-35.
14. Akpan A, Morgan R. Oral candidiasis. *Postgrad Med J*. 2002 Aug;78(922):455-9.
15. Webb AK, Woolnough E. *Candida albicans* infection in adults with cystic fibrosis. *J R Soc Med*. 2006;99 Suppl 46:13-6.
16. Sawyer SM, Bowes G, Phelan PD. Vulvovaginal candidiasis in young women with cystic fibrosis. *BMJ*. 1994 Jun 18;308(6944):1609.
17. Lyon A GE, Bilton D. Is genital candida infection a significant problem for adults with cystic fibrosis. *J Cyst Fibros*. 2004;3:S99.
18. Woolnough EM SC, Dodd ME, Jones A, Webb AK. Is candidiasis a problem in adults with cystic fibrosis? A prospective study. *Thorax*. 2004;59:S78.
19. Nurbhai M, Grimshaw J, Watson M, Bond C, Mollison J, Ludbrook A. Oral versus intra-vaginal imidazole and triazole anti-fungal treatment of uncomplicated vulvovaginal candidiasis (thrush). *Cochrane Database Syst Rev*. 2007(4):CD002845.
20. Orens JB, Estenne M, Arcasoy S, Conte JV, Corris P, Egan JJ, et al. International guidelines for the selection of lung transplant candidates: 2006 update--a consensus report

1
2
3 from the Pulmonary Scientific Council of the International Society for Heart and Lung
4 Transplantation. *J Heart Lung Transplant*. 2006 Jul;25(7):745-55.
5
6

7
8 21. Helmi M, Love RB, Welter D, Cornwell RD, Meyer KC. Aspergillus infection in lung
9 transplant recipients with cystic fibrosis: risk factors and outcomes comparison to other types
10 of transplant recipients. *Chest*. 2003 Mar;123(3):800-8.
11
12

13
14 22. Schaenman JM, Rosso F, Austin JM, Baron EJ, Gamberg P, Miller J, et al. Trends in
15 invasive disease due to *Candida* species following heart and lung transplantation. *Transpl*
16 *Infect Dis*. 2009 Apr;11(2):112-21.
17
18

19
20 23. Grossi P, Farina C, Fiocchi R, Dalla Gasperina D. Prevalence and outcome of
21 invasive fungal infections in 1,963 thoracic organ transplant recipients: a multicenter
22 retrospective study. Italian Study Group of Fungal Infections in Thoracic Organ Transplant
23 Recipients. *Transplantation*. 2000 Jul 15;70(1):112-6.
24
25
26

27
28 24. Bonacorsi SP, Munck A, Gerardin M, Doit C, Brahimi N, Navarro J, et al. In situ
29 management and molecular analysis of candidaemia related to a totally implantable vascular
30 access in a cystic fibrosis patient. *J Infect*. 1996 Jul;33(1):49-51.
31
32

33
34 25. Horn CK, Conway SP. Candidaemia: risk factors in patients with cystic fibrosis who
35 have totally implantable venous access systems. *J Infect*. 1993 Mar;26(2):127-32.
36
37

38
39 26. Munck A, Malbezin S, Bloch J, Gerardin M, Lebourgeois M, Derelle J, et al. Follow-
40 up of 452 totally implantable vascular devices in cystic fibrosis patients. *Eur Respir J*. 2004
41 Mar;23(3):430-4.
42
43

44
45 27. Douglas LJ. Medical importance of biofilms in *Candida* infections. *Rev Iberoam*
46 *Micol*. 2002 Sep;19(3):139-43.
47
48

49
50 28. Muthig M, Hebestreit A, Ziegler U, Seidler M, Muller FM. Persistence of *Candida*
51 species in the respiratory tract of cystic fibrosis patients. *Med Mycol*. 2009 Jan 31:1-8.
52
53
54
55
56
57
58
59
60

- 1
2
3 29. Navarro J, Rainisio M, Harms HK, Hodson ME, Koch C, Mastella G, et al. Factors
4 associated with poor pulmonary function: cross-sectional analysis of data from the ERCF.
5 European Epidemiologic Registry of Cystic Fibrosis. *Eur Respir J*. 2001 Aug;18(2):298-305.
6
7
8
9
10 30. Sullivan DJ, Westerneng TJ, Haynes KA, Bennett DE, Coleman DC. *Candida*
11 *dubliniensis* sp. nov.: phenotypic and molecular characterization of a novel species associated
12 with oral candidosis in HIV-infected individuals. *Microbiology*. 1995 Jul;141 (Pt 7):1507-
13 21.
14
15
16
17
18
19 31. Polacheck I, Strahilevitz J, Sullivan D, Donnelly S, Salkin IF, Coleman DC. Recovery
20 of *Candida dubliniensis* from non-human immunodeficiency virus-infected patients in Israel.
21 *J Clin Microbiol*. 2000 Jan;38(1):170-4.
22
23
24
25
26 32. Peltroche-Llacsahuanga H, Dohmen H, Haase G. Recovery of *Candida dubliniensis*
27 from sputum of cystic fibrosis patients. *Mycoses*. 2002 Feb;45(1-2):15-8.
28
29
30 33. Asmundsdottir LR, Erlendsdottir H, Agnarsson BA, Gottfredsson M. The importance
31 of strain variation in virulence of *Candida dubliniensis* and *Candida albicans*: results of a
32 blinded histopathological study of invasive candidiasis. *Clin Microbiol Infect*. 2009
33 Jun;15(6):576-85.
34
35
36
37
38
39 34. Hazen KC, Wu JG, Masuoka J. Comparison of the hydrophobic properties of *Candida*
40 *albicans* and *Candida dubliniensis*. *Infect Immun*. 2001 Feb;69(2):779-86.
41
42
43
44
45 35. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of
46 persistent infections. *Science*. 1999 May 21;284(5418):1318-22.
47
48
49
50 36. Hogan DA. Talking to themselves: autoregulation and quorum sensing in fungi.
51 *Eukaryot Cell*. 2006 Apr;5(4):613-9.
52
53
54 37. Williams P. Quorum sensing, communication and cross-kingdom signalling in the
55 bacterial world. *Microbiology*. 2007 Dec;153(Pt 12):3923-38.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
38. Hogan DA, Kolter R. *Pseudomonas-Candida* interactions: an ecological role for virulence factors. *Science*. 2002 Jun 21;296(5576):2229-32.
39. Joint I, Tait K, Callow ME, Callow JA, Milton D, Williams P, et al. Cell-to-cell communication across the prokaryote-eukaryote boundary. *Science*. 2002 Nov 8;298(5596):1207.
40. Hughes WT, Kim HK. Mycoflora in cystic fibrosis: some ecologic aspects of *Pseudomonas aeruginosa* and *Candida albicans*. *Mycopathol Mycol Appl*. 1973 Jul 31;50(3):261-9.
41. Williams P, Camara M. Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Curr Opin Microbiol*. 2009 Apr;12(2):182-91.
42. Lo HJ, Kohler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. Nonfilamentous *C. albicans* mutants are avirulent. *Cell*. 1997 Sep 5;90(5):939-49.
43. Hendrickson EL, Plotnikova J, Mahajan-Miklos S, Rahme LG, Ausubel FM. Differential roles of the *Pseudomonas aeruginosa* PA14 rpoN gene in pathogenicity in plants, nematodes, insects, and mice. *J Bacteriol*. 2001 Dec;183(24):7126-34.
44. Hornby JM, Jensen EC, Lisec AD, Tasto JJ, Jahnke B, Shoemaker R, et al. Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. *Appl Environ Microbiol*. 2001 Jul;67(7):2982-92.
45. Ramage G, Saville SP, Wickes BL, Lopez-Ribot JL. Inhibition of *Candida albicans* biofilm formation by farnesol, a quorum-sensing molecule. *Appl Environ Microbiol*. 2002 Nov;68(11):5459-63.
46. Cugini C, Calfee MW, Farrow JM, 3rd, Morales DK, Pesci EC, Hogan DA. Farnesol, a common sesquiterpene, inhibits PQS production in *Pseudomonas aeruginosa*. *Mol Microbiol*. 2007 Aug;65(4):896-906.

- 1
2
3
4 47. McAlester G, O'Gara F, Morrissey JP. Signal-mediated interactions between
5
6 Pseudomonas aeruginosa and Candida albicans. *J Med Microbiol*. 2008 May;57(Pt 5):563-9.
7
8 48. Dudler R, Eberl L. Interactions between bacteria and eukaryotes via small molecules.
9
10 *Curr Opin Biotechnol*. 2006 Jun;17(3):268-73.
11
12 49. Kobayashi H, Kobayashi O, Kawai S. Pathogenesis and clinical manifestations of
13
14 chronic colonization by Pseudomonas aeruginosa and its biofilms in the airway tract. *J Infect*
15
16 *Chemother*. 2009 Jun;15(3):125-42.
17
18 50. Jenkinson HF DL. Candida interactions with bacterial biofilms. Brogden KA GJ,
19
20 editor. Washington DC: ASM Press; 2002.
21
22 51. Rickard AH, Gilbert P, High NJ, Kolenbrander PE, Handley PS. Bacterial
23
24 coaggregation: an integral process in the development of multi-species biofilms. *Trends*
25
26 *Microbiol*. 2003 Feb;11(2):94-100.
27
28 52. El-Azizi MA, Starks SE, Khardori N. Interactions of Candida albicans with other
29
30 Candida spp. and bacteria in the biofilms. *J Appl Microbiol*. 2004;96(5):1067-73.
31
32 53. Tawara Y, Honma K, Naito Y. Methicillin-resistant Staphylococcus aureus and
33
34 Candida albicans on denture surfaces. *Bull Tokyo Dent Coll*. 1996 Aug;37(3):119-28.
35
36 54. Adam B, Baillie GS, Douglas LJ. Mixed species biofilms of Candida albicans and
37
38 Staphylococcus epidermidis. *J Med Microbiol*. 2002 Apr;51(4):344-9.
39
40 55. Baena-Monroy T, Moreno-Maldonado V, Franco-Martinez F, Aldape-Barrios B,
41
42 Quindos G, Sanchez-Vargas LO. Candida albicans, Staphylococcus aureus and Streptococcus
43
44 mutans colonization in patients wearing dental prosthesis. *Med Oral Patol Oral Cir Bucal*.
45
46 2005;10 Suppl 1:E27-39.
47
48 56. Carlson E. Effect of strain of Staphylococcus aureus on synergism with Candida
49
50 albicans resulting in mouse mortality and morbidity. *Infect Immun*. 1983 Oct;42(1):285-92.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
57. Carlson E. Enhancement by *Candida albicans* of *Staphylococcus aureus*, *Serratia marcescens*, and *Streptococcus faecalis* in the establishment of infection in mice. *Infect Immun.* 1983 Jan;39(1):193-7.
58. Jabra-Rizk MA, Meiller TF, James CE, Shirtliff ME. Effect of farnesol on *Staphylococcus aureus* biofilm formation and antimicrobial susceptibility. *Antimicrob Agents Chemother.* 2006 Apr;50(4):1463-9.
59. Levison ME, Pitsakis PG. Susceptibility to experimental *Candida albicans* urinary tract infection in the rat. *J Infect Dis.* 1987 May;155(5):841-6.
60. Kennedy MJ, Volz PA. Effect of various antibiotics on gastrointestinal colonization and dissemination by *Candida albicans*. *Sabouraudia.* 1985 Aug;23(4):265-73.
61. Holmes AR, Cannon RD, Jenkinson HF. Interactions of *Candida albicans* with bacteria and salivary molecules in oral biofilms. *J Ind Microbiol.* 1995 Sep;15(3):208-13.
62. O'Sullivan JM, Jenkinson HF, Cannon RD. Adhesion of *Candida albicans* to oral streptococci is promoted by selective adsorption of salivary proteins to the streptococcal cell surface. *Microbiology.* 2000 Jan;146 (Pt 1):41-8.
63. Cannon RD, Chaffin WL. Colonization is a crucial factor in oral candidiasis. *J Dent Educ.* 2001 Aug;65(8):785-7.
64. Holmes AR, van der Wielen P, Cannon RD, Ruske D, Dawes P. *Candida albicans* binds to saliva proteins selectively adsorbed to silicone. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006 Oct;102(4):488-94.
65. Basson NJ. Competition for glucose between *Candida albicans* and oral bacteria grown in mixed culture in a chemostat. *J Med Microbiol.* 2000 Nov;49(11):969-75.
66. Peleg AY, Tampakakis E, Fuchs BB, Eliopoulos GM, Moellering RC, Jr., Mylonakis E. Prokaryote-eukaryote interactions identified by using *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A.* 2008 Sep 23;105(38):14585-90.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
67. Ramage G, Tomsett K, Wickes BL, Lopez-Ribot JL, Redding SW. Denture stomatitis: a role for *Candida* biofilms. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004 Jul;98(1):53-9.
68. Lussier M, Sdicu AM, Shahinian S, Bussey H. The *Candida albicans* KRE9 gene is required for cell wall beta-1, 6-glucan synthesis and is essential for growth on glucose. *Proc Natl Acad Sci U S A.* 1998 Aug 18;95(17):9825-30.
69. Buurman ET, Westwater C, Hube B, Brown AJ, Odds FC, Gow NA. Molecular analysis of CaMnt1p, a mannosyl transferase important for adhesion and virulence of *Candida albicans*. *Proc Natl Acad Sci U S A.* 1998 Jun 23;95(13):7670-5.
70. Bulawa CE, Miller DW, Henry LK, Becker JM. Attenuated virulence of chitin-deficient mutants of *Candida albicans*. *Proc Natl Acad Sci U S A.* 1995 Nov 7;92(23):10570-4.
71. Chauhan N, Li, D., Singh, P., Calderone, R., Kruppa, M. . The Cell wall of *Candida* spp. Calderone RA, editor. Washington DC: ASM Press; 2002.
72. Masuoka J. Surface glycans of *Candida albicans* and other pathogenic fungi: physiological roles, clinical uses, and experimental challenges. *Clin Microbiol Rev.* 2004 Apr;17(2):281-310.
73. Timpel C, Strahl-Bolsinger S, Ziegelbauer K, Ernst JF. Multiple functions of Pmt1p-mediated protein O-mannosylation in the fungal pathogen *Candida albicans*. *J Biol Chem.* 1998 Aug 14;273(33):20837-46.
74. Csank C, Schroppel K, Leberer E, Harcus D, Mohamed O, Meloche S, et al. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infect Immun.* 1998 Jun;66(6):2713-21.
75. Cutler JE. N-glycosylation of yeast, with emphasis on *Candida albicans*. *Med Mycol.* 2001;39 Suppl 1:75-86.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
76. Ernst JF, Prill SK. O-glycosylation. *Med Mycol.* 2001;39 Suppl 1:67-74.
77. Kilpatrick DC. Mannan-binding lectin: clinical significance and applications. *Biochim Biophys Acta.* 2002 Sep 19;1572(2-3):401-13.
78. Brouwer N, Dolman KM, van Houdt M, Sta M, Roos D, Kuijpers TW. Mannose-binding lectin (MBL) facilitates opsonophagocytosis of yeasts but not of bacteria despite MBL binding. *J Immunol.* 2008 Mar 15;180(6):4124-32.
79. Lee SJ, Gonzalez-Aseguinolaza G, Nussenzweig MC. Disseminated candidiasis and hepatic malarial infection in mannose-binding-lectin-A-deficient mice. *Mol Cell Biol.* 2002 Dec;22(23):8199-203.
80. Brown GD. Dectin-1: a signalling non-TLR pattern-recognition receptor. *Nat Rev Immunol.* 2006 Jan;6(1):33-43.
81. Gantner BN, Simmons RM, Canavera SJ, Akira S, Underhill DM. Collaborative induction of inflammatory responses by dectin-1 and Toll-like receptor 2. *J Exp Med.* 2003 May 5;197(9):1107-17.
82. Brown GD, Herre J, Williams DL, Willment JA, Marshall AS, Gordon S. Dectin-1 mediates the biological effects of beta-glucans. *J Exp Med.* 2003 May 5;197(9):1119-24.
83. McGreal EP, Rosas M, Brown GD, Zamze S, Wong SY, Gordon S, et al. The carbohydrate-recognition domain of Dectin-2 is a C-type lectin with specificity for high mannose. *Glycobiology.* 2006 May;16(5):422-30.
84. Sato K, Yang XL, Yudate T, Chung JS, Wu J, Luby-Phelps K, et al. Dectin-2 is a pattern recognition receptor for fungi that couples with the Fc receptor gamma chain to induce innate immune responses. *J Biol Chem.* 2006 Dec 15;281(50):38854-66.
85. Jouault T, El Abed-El Behi M, Martinez-Esparza M, Breuilh L, Trinel PA, Chamaillard M, et al. Specific recognition of *Candida albicans* by macrophages requires

- 1
2
3
4 galectin-3 to discriminate *Saccharomyces cerevisiae* and needs association with TLR2 for
5
6 signaling. *J Immunol.* 2006 Oct 1;177(7):4679-87.
7
8 86. Cambi A, Gijzen K, de Vries JM, Torensma R, Joosten B, Adema GJ, et al. The C-
9
10 type lectin DC-SIGN (CD209) is an antigen-uptake receptor for *Candida albicans* on
11
12 dendritic cells. *Eur J Immunol.* 2003 Feb;33(2):532-8.
13
14
15 87. Kery V, Krepinsky JJ, Warren CD, Capek P, Stahl PD. Ligand recognition by purified
16
17 human mannose receptor. *Arch Biochem Biophys.* 1992 Oct;298(1):49-55.
18
19
20 88. Lee TW, Brownlee KG, Conway SP, Denton M, Littlewood JM. Evaluation of a new
21
22 definition for chronic *Pseudomonas aeruginosa* infection in cystic fibrosis patients. *J Cyst*
23
24 *Fibros.* 2003 Mar;2(1):29-34.
25
26
27 89. Wells CA, Salvage-Jones JA, Li X, Hitchens K, Butcher S, Murray RZ, et al. The
28
29 macrophage-inducible C-type lectin, mincle, is an essential component of the innate immune
30
31 response to *Candida albicans*. *J Immunol.* 2008 Jun 1;180(11):7404-13.
32
33
34 90. Lemaitre B, Nicolas E, Michaut L, Reichhart JM, Hoffmann JA. The dorsoventral
35
36 regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in
37
38 *Drosophila* adults. *Cell.* 1996 Sep 20;86(6):973-83.
39
40
41 91. Simon A, Kullberg BJ, Tripet B, Boerman OC, Zeeuwen P, van der Ven-Jongekrijg J,
42
43 et al. Drosomycin-like defensin, a human homologue of *Drosophila melanogaster* drosomycin
44
45 with antifungal activity. *Antimicrob Agents Chemother.* 2008 Apr;52(4):1407-12.
46
47
48 92. Ozinsky A, Underhill DM, Fontenot JD, Hajjar AM, Smith KD, Wilson CB, et al. The
49
50 repertoire for pattern recognition of pathogens by the innate immune system is defined by
51
52 cooperation between toll-like receptors. *Proc Natl Acad Sci U S A.* 2000 Dec
53
54 5;97(25):13766-71.
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
93. Netea MG, Van Der Graaf CA, Vonk AG, Verschueren I, Van Der Meer JW, Kullberg BJ. The role of toll-like receptor (TLR) 2 and TLR4 in the host defense against disseminated candidiasis. *J Infect Dis.* 2002 May 15;185(10):1483-9.
94. Villamon E, Gozalbo D, Roig P, O'Connor JE, Fradelizi D, Gil ML. Toll-like receptor-2 is essential in murine defenses against *Candida albicans* infections. *Microbes Infect.* 2004 Jan;6(1):1-7.
95. Netea MG, van de Veerdonk F, Verschueren I, van der Meer JW, Kullberg BJ. Role of TLR1 and TLR6 in the host defense against disseminated candidiasis. *FEMS Immunol Med Microbiol.* 2008 Jan;52(1):118-23.
96. Blasi E, Mucci A, Neglia R, Pezzini F, Colombari B, Radzioch D, et al. Biological importance of the two Toll-like receptors, TLR2 and TLR4, in macrophage response to infection with *Candida albicans*. *FEMS Immunol Med Microbiol.* 2005 Apr 1;44(1):69-79.
97. Netea MG, Suttmuller R, Hermann C, Van der Graaf CA, Van der Meer JW, van Krieken JH, et al. Toll-like receptor 2 suppresses immunity against *Candida albicans* through induction of IL-10 and regulatory T cells. *J Immunol.* 2004 Mar 15;172(6):3712-8.
98. Suttmuller RP, den Brok MH, Kramer M, Bennink EJ, Toonen LW, Kullberg BJ, et al. Toll-like receptor 2 controls expansion and function of regulatory T cells. *J Clin Invest.* 2006 Feb;116(2):485-94.
99. van de Veerdonk FL, Netea MG, Jansen TJ, Jacobs L, Verschueren I, van der Meer JW, et al. Redundant role of TLR9 for anti-*Candida* host defense. *Immunobiology.* 2008;213(8):613-20.
100. Greene CM, McElvaney NG. Toll-like receptor expression and function in airway epithelial cells. *Arch Immunol Ther Exp (Warsz).* 2005 Sep-Oct;53(5):418-27.
101. Greene CM, Branagan P, McElvaney NG. Toll-like receptors as therapeutic targets in cystic fibrosis. *Expert Opin Ther Targets.* 2008 Dec;12(12):1481-95.

- 1
2
3 102. Mencacci A, Cenci E, Del Sero G, Fe d'Ostiani C, Mosci P, Trinchieri G, et al. IL-10
4 is required for development of protective Th1 responses in IL-12-deficient mice upon
5
6
7
8
9
10
11 103. Vazquez-Torres A, Jones-Carson J, Wagner RD, Warner T, Balish E. Early resistance
12 of interleukin-10 knockout mice to acute systemic candidiasis. *Infect Immun*. 1999
13
14
15
16
17
18 104. Lavigne LM, Schopf LR, Chung CL, Maylor R, Sypek JP. The role of recombinant
19 murine IL-12 and IFN-gamma in the pathogenesis of a murine systemic *Candida albicans*
20
21
22
23
24
25 105. Allard JB, Rinaldi L, Wargo MJ, Allen G, Akira S, Uematsu S, et al. Th2 allergic
26 immune response to inhaled fungal antigens is modulated by TLR-4-independent bacterial
27
28
29
30
31
32 106. Roilides E, Holmes A, Blake C, Pizzo PA, Walsh TJ. Effects of granulocyte colony-
33 stimulating factor and interferon-gamma on antifungal activity of human polymorphonuclear
34
35
36
37
38
39
40
41 107. Gaviria JM, van Burik JA, Dale DC, Root RK, Liles WC. Modulation of neutrophil-
42 mediated activity against the pseudohyphal form of *Candida albicans* by granulocyte colony-
43
44
45
46
47
48 108. Przyklenk B, Bauernfeind A, Horl G, Emminger G. Serologic response to *Candida*
49
50
51
52
53 109. Maiz L, Cuevas M, Quirce S, Canon JF, Pacheco A, Sousa A, et al. Serologic IgE
54
55
56
57
58
59
60

1
2
3
4 110. Maiz L, Cuevas M, Lamas A, Sousa A, Quirce S, Suarez L. [Aspergillus fumigatus
5
6 and Candida albicans in cystic fibrosis: clinical significance and specific immune response
7
8 involving serum immunoglobulins G, A, and M]. Arch Bronconeumol. 2008 Mar;44(3):146-
9
10 51.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only